

About the quark currents in leptonic decays of pseudoscalar mesons

V.P Efrosinin, A.N. Khotjantsev

Institute for Nuclear Research RAS

60th October Ave., 7A, 117312, Moscow, Russia

Abstract

With usage of obvious mechanisms of quark currents the amplitudes of leptonic decays of pseudoscalar mesons are investigated. The estimation for a constant of leptonic decay of the D^+ meson is obtained, $f_{D^+} \simeq 0.23 \text{ GeV}$.

In the paper [1] by the calculation of form factors in $K_{\mu 3}$ decay in spectator nearing the research of obvious mechanisms of quark currents was conducted. Thus the output for frameworks of the conventional phenomenological description of weak decays of hadrons was carried out. In the present paper are investigated quark currents in leptonic decays of pseudoscalar mesons.

Let us consider $\pi^+ \rightarrow \mu^+ \nu_\mu$ decay (Fig. 1). The amplitude of the process has a kind:

$$M = \frac{G_F}{\sqrt{2}} V_{ud}^* \bar{u}_\nu \gamma^\alpha (1 - \gamma_5) v_\mu < 0 | J_\alpha | \pi^+ >, \quad (1)$$

where the phenomenological matrix element of a pseudovector current can be submitted as

$$< 0 | J_\alpha | \pi^+ > = \varphi_\pi f_\pi p_\alpha. \quad (2)$$

Here f_π is a constant of a charged pion, $f_\pi = 0.131 \text{ GeV}$ [2], φ_π is a wave function of a pion, and p_α is its four-momentum.

As well as in paper [1] we record a pseudovector current as a current of constituent quarks

$$< 0 | J_\alpha | \pi^+ > = \bar{v}_{\bar{d}} \gamma_\alpha \gamma_5 u_u F_\pi, \quad (3)$$

here $\bar{v}_{\bar{d}}, u_u$ are four-spinors of constituent quarks, F_π is parameter of an annihilation in vacuum, having dimension of mass. F_π corresponds to an annihilation of a

virtual pair of $u\bar{u}$. And the degree of this process (departure from a mass surface) corresponds to energy of a pair lepton-antilepton or mass of a pion.

In the rest frame of a pion from (3) by simple calculation is received

$$< 0 | J_\alpha | \pi^+ > = F_\pi 2m_u, \quad (4)$$

where m_u is mass of a constituent $u(d)$ quark, as well as in the paper [1] and also in the papers [3, 4]. From expressions (2) and (4) is received

$$f_\pi m_\pi = F_\pi 2m_u, \quad (5)$$

$$f_\pi = 0.131 \text{ GeV} [2].$$

Similarly for $K^+ \rightarrow \mu^+ \nu_\mu$ decay we have

$$f_K m_K = F_K 2\sqrt{m_u m_s}, \quad (6)$$

$$f_K = 0.160 \text{ GeV} [2].$$

And also for $D_s^+ \rightarrow \mu^+ \nu_\mu$ decay is

$$f_{D_s^+} m_{D_s^+} = F_{D_s^+} 2\sqrt{m_c m_s}, \quad (7)$$

$$f_{D_s^+} \simeq 0.288 \text{ GeV} [2].$$

Using (5,6,7) and also masses of constituent quarks

$$\begin{aligned} m_u &= 0.305 \text{ GeV} [1], \\ m_s &= 0.487 \text{ GeV} [1], \\ m_c &= 1.400 \text{ GeV} [2] \end{aligned}$$

is received

$$\begin{aligned} F_{\pi^+} &= \frac{f_\pi m_\pi}{2m_u} = 0.030 \text{ GeV}, \\ F_{K^+} &= \frac{f_K m_K}{2\sqrt{m_u m_s}} = 0.103 \text{ GeV}, \\ F_{D_s^+} &= \frac{f_{D_s^+} m_{D_s^+}}{2\sqrt{m_c m_s}} = 0.343 \text{ GeV}. \end{aligned} \quad (8)$$

We are interested by a prediction for a constant of $D^+ \rightarrow \mu^+ \nu$ decay. In paper [2] for parameter f_{D^+} the high bound is given only

$$f_{D^+} < 310 \text{ MeV}. \quad (9)$$

We have suspected that parameter F_{M^+} depends only on mass of a meson. We search for this relation as

$$F_{M^+} = aM^2 + bM + c. \quad (10)$$

With usage of (8) we discover

$$F_{M^+} = -0.0238M^2 + 0.2213M - 0.0005 \text{ (GeV)}. \quad (11)$$

The behavior of parameter F_{M^+} depending on mass is submitted in (Fig. 2).

Let us remark that a matrix element in (3) is half outside of a mass surface. And is approximated as on mass surface. The effect of departure from a mass surface is actuated in parameter of an annihilation F_{M^+} . The difference of these effects for pairs $u\bar{u}$ and $s\bar{s}$ is supposed minor.

From the formula (11) is received

$$F_{D^+} = 0.33 \text{ GeV}. \quad (12)$$

Then from expression

$$F_{D^+} m_{G^+} = F_{D^+} 2\sqrt{m_c m_u} \quad (13)$$

we have

$$f_{D^+} = 0.23 \text{ GeV}, \quad (14)$$

that does not contradict (9).

The obtaining of the new data of constants of decay of pseudoscalar mesons will help to explain a physical essence of parameter of an annihilation F_{M^+} . The introducing of this parameter encourages division of series stages weak and strong interactions in decays of pseudoscalar mesons and refinement of mechanism of these decays.

References

- [1] V. P. Efrosinin and A. N. Khotyantsev, arXiv:hep-ph/0510169.
- [2] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592** (2004) 1.
- [3] S. S. Gershtein and M. Y. Khlopov, Pisma Zh. Eksp. Teor. Fiz. **23** (1976) 374.
- [4] M. Y. Khlopov, Yad. Fiz. **18** (1978) 1134.




Figure 1:




Figure 2:

Figure captions

Fig. 1. The diagram of $\pi^+ \rightarrow \mu^+ \nu_\mu$ decay on a quark level.

Fig. 2. Relation of parameter of an annihilation F_{M^+} to mass of a pseudoscalar meson.